arXiv:1209.0209v1 [hep-th] 2 Sep 2012

Lorentz invariance of vacuum in the closed time path formalism

Adam Bednorz[1, *]

[1]*Faculty of Physics, University of Warsaw ul. Hoża 69, PL00-681 Warsaw, Poland*

(Dated: September 4, 2012)

The Schwinger-Kadanoff-Baym-Keldysh closed time path formalism is fundamentally important because it combines the quantum statistical approach with arbitrary field theories. We prove that in this approach the Green functions in vacuum at zero temperature and vanishing conserved charges are invariant with respect to continuous Lorentz transformations. To make the proof clear we adopt the well-known techniques of path integrals and Feynman diagrams tailored especially to the closed time path by full use of the properties of complex time. The proof, which is in principle perturbative, relies on the quantum electrodynamics Lagrangian and extends to the Standard Model in general.

I. INTRODUCTION

Quantum field theory (QFT) is the basis of all modern physical theories [1, 2], including quantum electrodynamics, the Standard Model, and all their approximations, in particular nonrelativistic, mean-field and various condensed-matter models. Despite the same root – the historical problem of black-body radiation – quantum particle physics and quantum statistical physics underwent somewhat separate developments, although in a related way. The former lead e.g. to the beautiful Feynmann diagrams in spacetime [3] while the latter required making the time imaginary, to obtain temperature [4, 5]. The road to reunify dynamics and statistical physics was not straightforward. Schwinger, Kadanoff, Baym, Keldysh, [6–8] and others [9, 10] have constructed the closed time path (CTP) formalism, in which the time is complex and can represent both the usual time (real axis) and the temperature (imaginary axis). The time on the real axis flows first forwards and then backwards. This happens also in zero-temperature vacuum QFT, where the scattering cross sections are given by the modulus square of the amplitude. The amplitude and its complex conjugate correspond to the forward and backward part of the time path, respectively. They are connected via Cutkosky rules [11].

The CTP formalism provides a scheme to calculate correlation functions of physical quantities along the time path, defining a family of Green functions [12–14]. Due to analyticity of time functions, there are numerous relations between Green functions [12, 15], constituting generalizations of fluctuation-dissipation theorem (FDT) [16]. The range of applications of CTP is very wide, from non-equlibrium in condensed matter systems [17] through quantum optics [18] to fundamental theories and cosmology [19, 20]. Since only initial conditions are required CTP is sometimes called in-in formalism to distinguish it from in-out scattering picture. In static problems in equilibrium one can omit the real part of the path and work with Matsubara frequencies, corresponding to the Fourier transform of the imaginary part of the path. The Matsubara formalism is widely used to determine thermodynamic properties and phase transitions of many-body systems [4, 5].

The formalism works fine both in operator and field-path-integral way [21]. However, the Lorentz invariance of vacuum (ground state) Green functions at zero temperature and zero average charges is not manifest because of the preferred time coordinate to form the complex contour. To our best knowledge, it is believed only intuitively [14, 22], while the strict proof has never been demonstrated. For a comparison, Lorentz invariance of individual fields has been shown by Schwinger [23] while the invariance of the in-out scattering matrix is textbook knowledge [2]. However, for the CTP formalism, one needs Lorentz invariance of the full vacuum ground state and arbitrary in-in fields. An attempt to prove Lorentz invariance by means of energy-momentum representation and trying to reduce the proof to familiar in-out amplitudes raises further problems instead, e.g. pinch singularities [24, 25], contribution of imaginary times to the self-energy [26, 27] and non-factorization of generating functional [14, 28, 29]. Using momentum representation one can only prove the Lorentz invariance of particular classes of diagrams (e.g. lowest-order self-energy of elementary particles) but not in general. The Lorentz invariance could be certainly proved only if the underlying Lagrangian is Lorentz invariant and conserved charges are zero. It has been suggested that Nature may violate the invariance directly (by non-invariant Lagrangian) or spontaneously (by a Higgs-like mechanism) [30]. We do not discuss these two last possibilities here.

In this paper, we confirm the common intuition by a direct proof. Although formally general, it is better to understand it as perturbative. The CTP formalism can be indeed formulated both in Hamiltonian and Lagrangian way under several conditions, satisfied by all known QFTs. The zero-temperature vacuum Green functions are invariant under continuous Lorentz transformations. However, the discrete transformations like time-reversal and parity have to be treated separately. If the underlying Lagrangian is invariant under charge conjugation or parity reflection then the proof remains valid for these transformations [31]. Unfortunately, even if the

---

[*] Electronic address: Adam.Bednorz@fuw.edu.pl

Lagrangian is symmetric with respect to time reversal, then CTP is not (except special cases, e.g. at space-like points or in scattering problems). In the same way, CTP is in general not invariant under joint charge-parity-time reversal in contrast to Lagrangian.

The paper is organized as follows. We firs recall the construction of the complex time Hamiltonian and Lagrangian CTP, defining relevant Green functions, leaving details on Appendices. These parts are not much original, but adopted from various textbooks and papers to CTP – with the essential generalization to complex times. Then we prove the Lorentz invariance of the zero-temperature no-charge vacuum for continuous transformations, which is the main original result of this work. Finally, we discuss possible pitfalls in a approaches to a proof based on momentum representation and the problem of time asymmetry. Throughout the paper, we use the convention $\hbar = c = 1$ and $\beta = 1/k_B T$.

II. CTP IN THE HAMILTONIAN PICTURE

Let us begin with the textbook picture of CTP. The starting point of every quantum-mechanical problem is the Hermitian Hamiltonian operator $\hat{H}$ in a given Hilbert space. The physical quantities are described by Hermitian operators $\hat{X}_k$ corresponding to their classical counterparts X_k (numbers). Depending on the measurement scheme, there exists a correspondence between operators and measurement outcomes, $\hat{X}_k \to X_k$. The system is described by its initial state, given by its density matrix $\hat{\rho}$, which is Hermitian, positive definite and normalized, $\mathrm{Tr}\hat{\rho} = 1$. It can be always expressed in the form $\hat{\rho} = \sum_k p_k |k\rangle\langle k|$, where the states $|k\rangle$ form a complete, orthonormal basis in the Hilbert space, $\langle k|q\rangle = \delta_{kq}$, and $\sum_k p_k = 1$, $p_k \geq 0$. Having the initial state, one can continuously evolve the state in time t according to the Hamiltonian $\hat{H}(t) \to \hat{H} + \hat{H}_1(t)$, where $\hat{H}_1(t)$ is the time dependent part, due to changing external forces, which are assumed to be absent for $t < 0$. For the main purpose of this paper, $\hat{H}_1$ can be omitted. However, the whole CTP formalism works perfectly also for nonzero $\hat{H}_1$, which is necessary in problems involving external (changing) forces. The projection postulate defines averages in the projective (von Neumann) detection scheme [32] for commuting observables, $\hat{X}_k\hat{X}_l = \hat{X}_l\hat{X}_k$,

$$\left\langle \prod_k X_k(t) \right\rangle_\rho := \mathrm{Tr}\hat{\rho} \times \left[\tilde{\mathcal{T}} e^{\int_0^t i\hat{H}(t')dt'}\right] \left[\prod_k \hat{X}_k\right] \left[\mathcal{T} e^{-\int_0^t i\hat{H}(t')dt'}\right], \quad (1)$$

where $\mathcal{T}(\tilde{\mathcal{T}})$ denotes (anti)time ordering of operators in the Taylor expansion, namely

$$\mathcal{T}\hat{Y}_1(t_1)\cdots\hat{Y}_n(t_n) = \mathrm{sgn}(\sigma_f)\hat{Y}_{\sigma(1)}(t_{\sigma(1)})\cdots\hat{Y}_{\sigma(n)}(t_{\sigma(n)}), \quad (2)$$



FIG. 1: The Keldysh contour. The path lies infinitesimally close to real axis and reverses its direction at some final time t_f

where σ is a permutation such that $t_{\sigma(1)} \geq \cdots \geq t_{\sigma(n)}$. Here $\mathrm{sgn}(\sigma_f)$ denotes the sign of the sub-permutation fermionic operators. If $\hat{Y}_k$ is a fermionic operator for $k \in \{1..n_f\}$, then σ_f is defined as the map between the ordered sequences $(\sigma(k))_{k=1}^{n_f}$ and $(\sigma(k))$, ordered according to k and $\sigma(k)$, respectively. For $\tilde{\mathcal{T}}$, we require $t_{\sigma(1)} \leq \cdots \leq t_{\sigma(n)}$. In the Heisenberg picture, it is convenient to define operators

$$\hat{A}_H(t) = \tilde{\mathcal{T}} e^{\int_0^t i\hat{H}(t')dt'} \hat{A} \mathcal{T} e^{-\int_0^t i\hat{H}(t')dt'}. \quad (3)$$

For practical applications of CTP like Feynman diagrams one is forced to define generalized averages [12], without referring to a particular detection scheme, constituting the family of Green functions,

$$\left\langle \prod_k X_k(t_k) \right\rangle_{q,\rho} = \mathrm{Tr}\hat{\rho}\tilde{\mathcal{T}} \prod_{k<q} \hat{X}_{kH}(t_k) \mathcal{T} \prod_{k>q} \hat{X}_{kH}(t_k). \quad (4)$$

Already at this point, it convenient to define the Keldysh contour [6, 8], where the time variable flows first from 0 to some final time t_0 (not necessarily infinite) and then goes backwards. In the complex plane, one can think of a (smooth) mapping $s \to t(s)$, where s is a strictly real variable ($s \in [0, 2t_0]$) with $t(s < t_0) = s + i\epsilon$, $t(s > t_0) = 2t_0 - s - i\epsilon$, such that the two branches are connected smoothly at t_0 and $\epsilon \to 0_+$, see Fig. 1 .

We introduce Keldysh ordering $\mathcal{T}_K$ defined as (2) but with $s_{\sigma(1)} \geq \cdots \geq s_{\sigma(n)}$. It combines both $\mathcal{T}$ (for $s < t_0$) and $\tilde{\mathcal{T}}$ (for $s > t_0$). We can write now

$$\left\langle \prod_k X_k(t_k) \right\rangle_{q,\rho} = \mathrm{Tr}\hat{\rho}\mathcal{T}_K \prod_k \hat{X}_{kH}(t(s_k)), \quad (5)$$

with $s_k \lesseqgtr t_0$ for $k \lesseqgtr q$, so that $t_k \equiv t(s_k)$ lies infinitesimally above/below the real axis. More generally, we can define the generating functional

$$\begin{aligned} e^{S[\chi]} &= \left\langle e^{\sum_k \int_K i\chi_k(t)X_k(t)dt} \right\rangle \\ &= \mathrm{Tr}\hat{\rho}\mathcal{T}_K e^{\sum_k \int_K i\chi_k(t)\hat{X}_{kH}(t)dt} \\ &= \mathrm{Tr}\hat{\rho}\mathcal{T}_K e^{\int_K idt\left(\sum_k \chi_k(t)\hat{X}_k - \hat{H}(t)\right)}. \end{aligned} \quad (6)$$

Here $dt = (dt/ds)ds$ and the integration runs over the whole Keldysh contour (denoted by subscript K), parametrized by the real parameter s. The averages (5) are given by derivatives of the generating functional,

$$\left\langle \prod_k X_k(t_k) \right\rangle_{q,\rho} = \prod_{k<q} \frac{-i\delta}{\delta\chi_k(t_k)} \prod_{k\geq q} \frac{i\delta}{\delta\chi_k(t_k)}\, e^{S[\chi]}\Big|_{\chi=0}. \tag{7}$$

After taking Keldysh ordering in (6), the information which Keldysh part $\chi(t_k)$ belongs to is encoded in the sign of the imaginary part of t_k. It is important that the averages do not depend on the turning time t_0 as long as it is later than all the (real parts of) arguments in the average. The Keldysh contour is also very useful in the interaction picture, where $\hat{H}(t) = \hat{H}_0 + \hat{H}_I(t)$ with the noninteracting part $\hat{H}_0$. In analogy to Heisenberg operators (3), we define interaction picture

$$\hat{X}_0(t) = e^{it\hat{H}_0}\hat{X}e^{-it\hat{H}_0}. \tag{8}$$

We can rewrite the generating functional in the form

$$e^{S[\chi]} = \mathrm{Tr}\hat{\rho}\mathcal{T}_K e^{\int_K idt\left(\sum_k \chi_k(t)\hat{X}_{k0}(t) - \hat{H}_{I0}(t)\right)}, \tag{9}$$

where both $\hat{A}_{k0}$ and $\hat{H}_{I0}$ are defined by (8). The formula (9) is very practical if the initial state can be expressed in terms of eigenstates of $\hat{H}_0$ as we can use perturbative methods to evaluate the desired Green functions.

It is often convenient to introduce four *superoperators* $\check{A}$, including the classical (Keldysh) and the quantum superoperator,

$$\begin{aligned}
&\check{A}^+\hat{\rho} = \hat{A}\hat{\rho},\\
&\check{A}^-\hat{\rho} = \hat{\rho}\hat{A},\\
&\check{A}^c = (\check{A}^+ + \check{A}^-)/2 = \{\hat{A},\cdot\}/2,\\
&\check{A}^q = \check{A}^+ - \check{A}^- = [\hat{A},\cdot],
\end{aligned} \tag{10}$$

where $[\hat{A},\hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}$ denotes the commutator and $\{\hat{A},\hat{B}\} = \hat{A}\hat{B} + \hat{B}\hat{A}$ the anticommutator. The superoperators are linear transformations on operators, $\check{X}\hat{O} = \hat{O}'$. It is straightforward to generalize the Hamiltonian picture to superoperators, with Heisenberg equation

$$\check{A}_H(t) = \tilde{\mathcal{T}} e^{\int_0^t i\check{H}^q(t')dt'}\check{A}\mathcal{T}e^{-\int_0^t i\check{H}^q(t')dt'}. \tag{11}$$

We define averages similarly to (5),

$$\left\langle \prod_k X^x_k(t_k) \right\rangle_\rho = \mathrm{Tr}\mathcal{T}\prod_k \check{X}^x_{kH}(t_k)\hat{\rho} \tag{12}$$

where $x = +,-,c,q$. Let us denote the two branches of Keldysh contour by $t_\pm = t \pm i\epsilon$. The averages (12) can be obtained from (5) by identifying $X^\pm(t) = X(t_\pm)$, $X^c = (X^+ + X^-)/2$ and $X^q = X^+ - X^-$. An important property of $X^q(t)$ is that if its time argument is the latest time with respect to other quantities then the average vanishes,

$$\langle \cdots X^q(t_{max}) \cdots \rangle_\rho = 0, \tag{13}$$

which follows from the cyclic property of the trace.

The initial state $\hat{\rho}$ is usually not arbitrary but derived from basic assumptions: stationarity, energy conservation, boundary conditions and other conservation laws. If the state is stationary then $[\hat{H},\hat{\rho}] = 0$. In equilibrium, the generally assumed structure is

$$\hat{\rho} \propto \exp\left[-\beta(\hat{H} - \mu_Q\hat{Q})\right] \tag{14}$$

where $\beta = 1/k_BT$ is the inverse temperature and $\hat{Q}$ is some conserved quantity (e.g. total charge, lepton or baryon number, depending on Lagrangian structure), $[\hat{Q},\hat{H}] = 0$ and μ_Q is the appropriate Lagrange multiplier. Every conserved quantity can be incorporated into the definition of the Hamiltonian, $\hat{H} \to \hat{H} - \mu_Q\hat{Q}$. Other stationary, but not equilibrium states may differ from (14) but they need a careful discussion of boundary conditions. In most applications one can think of initial thermal state (14), which is driven out from equilibrium by external forces, added via $\hat{H}_1(t)$, and one is interested in the response. Below, we assume (14) with $\mu_Q = 0$ (or equivalently $\hat{Q}$ is incorporated into $\hat{H}$).

Already from (14) it is clear that β can be understood as imaginary time. As suggested by Kadanoff and Baym [7], we extend the Keldysh contour to the smooth closed time path (CTP) in the complex plane of times, parametrized by real $s \in [s_i, s_f] \to t(s)$ with

$$\begin{aligned}
&\mathrm{Im}(dt/ds) \leq 0,\\
&t(s_f) = t(s_i) - i\beta,\\
&\mathrm{Im}\,t \to 0_+ \text{ and } \mathrm{Re}\,dt/ds > 0 \text{ for } s \in [s_+, s_0],\\
&\mathrm{Im}\,t \to 0_- \text{ and } \mathrm{Re}\,dt/ds < 0 \text{ for } s \in [s_0, s_-],\\
&t(s_\pm) = 0_\pm, t(s_0) = t_0.
\end{aligned} \tag{15}$$

Here $s_i < s_+ < s_0 < s_- < s_f$. The first condition is connected with the fact that usually $\hat{H}$ is bounded from below but not from above and we have to escape possible infinities. There is still an infinite freedom of possible choices of the path [10, 15, 33, 34]. One of possible paths is depicted in Fig. 2.

We define $\mathcal{T}_C$ by simple extending $\mathcal{T}_K$ to the whole CTP. The averages (12) can be written in the form

$$e^{S[\chi]} = \frac{\mathrm{Tr}\mathcal{T}_C e^{\int_C idt\left(\sum_k \chi_k(t)\hat{X}_k - \hat{H}(t)\right)}}{\mathrm{Tr}\mathcal{T}_C e^{-\int_C idt\hat{H}(t)}}, \tag{16}$$

where the subscript C denotes integration over the CTP. To facilitate an easier notation throughout the paper we define Heaviside and Dirac functions for complex time arguments t,

$$\begin{aligned}
&\theta(t-t') = \theta(s-s'),\ \delta(t-t') = \delta(s-s')/(dt/ds)\\
&\partial_t = (dt/ds)^{-1}\partial_s,\ dt = (dt/ds)ds
\end{aligned} \tag{17}$$

where as usual $\theta(s-s') = 1$ for $s > s'$ and 0 otherwise while $\int ds\delta(s-s') = 1$ and $\delta(s-s') = 0$ for $s \neq s'$. This is important since only s is the real, ordering parameter,



FIG. 2: The closed time path in the complex time plane. In the positive real part half-plane it is just the Keldysh contour from Fig. 1. The new parts lie in the negative real part half-plane and are connected smoothly to the Keldysh contour. All the path cannot go upwards in the imaginary direction. The finite temperature implies the jump $i\beta$, which extends to infinity at zero temperature. Apart from the above conditions, the shape of the path is arbitrary. We stress that the path is ordered by the real parameter s. All the important times have been marked.

while t is complex and thus has no natural ordering. Note that in all contour time-integrals we essentially integrate over s. Performing a Taylor expansion of exponentials in (16) one can explicitly check that the result does not depend on a particular shape of CTP [26], as long as (15) is satisfied and t_0 is greater than all interesting times. In such a check it is important to note that $\hat{H}$ is time-independent along the wiggly part of the contour, so the ordering does not matter there. On the other hand (16) is useful also for deriving the thermodynamic functions and transport coefficients, e.g. the Kubo formula [35]. In the interaction picture, we have to extend the definition (8) to arbitrary complex times. Then

$$e^{S[\chi]} = \frac{\mathrm{Tr}\hat{\rho}_0 \mathcal{T}_C e^{\int_C idt \left(\sum_k \chi_k(t)\hat{A}_{k0}(t) - \hat{H}_{I0}(t)\right)}}{\mathrm{Tr}\hat{\rho}_0 \mathcal{T}_C e^{-\int_C idt\, \hat{H}_{I0}(t)}}, \tag{18}$$

where $\hat{\rho}_0 \propto \exp(-\beta\hat{H}_0)$. The remaining calculations are usually performed perturbatively, with the help of the free two-point Green's functions and Wick's theorem [36, 37], which allows a decoupling of the free many-point Green functions into products of two-point functions, see Appendix A.

III. PATH INTEGRALS AND LAGRANGIAN

The Hamiltonian approach works as long as we do not care about relativity, because the Hamiltonian distinguishes a fixed reference frame. However, we would like to have a fully relativistic formalism. The way out is to turn to a Lagrangian, which leads to the path integral formulation of quantum field theory [3, 38, 39]. Here we adopt this well-known formulation to the CTP formalism. Although quite straightforward, it is very instructive to recall it in Appendix B(a-b), to see that CTP barely affects the path integral formalism. The only important point is that the time is complex, see (17).

It is convenient to formulate the generating functional for CTP averages with respect to the action, making use of the Lagrangian approach, namely

$$\left\langle \prod_k q_k(t_k) \prod_l \phi_l(t_l) \right\rangle = \prod_k \frac{\delta}{\delta i\chi_k(t_k)} \prod_l \frac{\delta}{\delta i\eta_l(t_l)} e^{S[\chi,\eta]}\Bigg|_{\chi=0,\eta=0} \tag{19}$$

where

$$e^{S[\chi,\eta]} = \frac{\int DAD\phi\, e^{\int_C idt(L+\sum_k(\chi_k(t)A_k(t)+\eta_k(t)\phi_k(t)))}}{\int DAD\phi\, e^{\int_C idt\, L}} \tag{20}$$

with auxiliary c-field χ and Grassmann field η. In the above, A and ϕ are generic bosonic and fermionic (Grassmann) fields, respectively, and the Lagrangian is a time-local function of fields and their time derivatives, namely $L = L[A, \dot{A}, \phi, \dot{\phi}]$.

The path-integral formulation of CTP is fully consistent with Wick theorem, charge conservation laws and renormalization, see Appendix B(c),(d) and (e), respectively, for details.

A. Contour shape independence

The fact that we can freely wiggle the contour is reasonably believed for CTP [10, 15, 33, 34], but a direct proof is very instructive [26]. Let us prove that (20) is independent of a particular shape of CTP, because the proof of Lorentz invariance will be analogous. It follows from the invariance of DA and $D\psi$ under infinitesimal transformation $\delta t(s)$ parametrized by real s (the variation may change endpoints but the jump must be kept constant)

$$\begin{aligned} A(t) &\to e^{\delta t\partial_t}A(t) = A(t) + \delta t\partial_t A(t) + \ldots \\ \phi(t) &\to e^{\delta t\partial_t}\phi(t) = \phi(t) + \delta t\partial_t \phi(t) + \ldots \end{aligned} \tag{21}$$

which gives $L(t+\delta t) = \exp(\delta t\partial_t)L(t)$. This means that the effect of the contour changes can be absorbed into a shift under time integral and a such it disappears (analogously to an integral over a total derivative). We stress that this proof is valid only if do not change the shape of the Keldysh part of the contour, except shifting t_0 if it lies beyond all relevant times. It is essential that the fields are smooth.

An alternative, more explicit proof relies on perturbative approach. One starts with Gaussian, diagonalized Lagrangian

$$\begin{aligned} L_0 \;=\; & \sum_k [\psi_k^*(i\partial_t - E_k)\psi_k \qquad (22) \\ & + \; A_k^*(i\partial_t - E_k)A_k + \dot{B}_k^2/2 - E_k^2 B_k^2/2)\Big] . \end{aligned}$$

In the above, we have denoted complex ψ instead of real ϕ and distinguished linear time-derivative bosonic field

A from quadratic B. The two-point Green function for B reads

$$\langle B_k(t)B_l(t')\rangle_0 = \frac{\delta_{kl}}{2E_k}\sum_{\pm}\mp\frac{e^{\pm i|t'-t|E_k}}{1-e^{\pm\beta E_k}}, \tag{23}$$

where $|t'-t| = (t-t')\theta(t-t') + (t'-t)\theta(t'-t)$. For A we get (A.4) and for ψ – (A.5).

If we add interaction term, $L = L_0 + L_I$, and L_I depends on fields appearing in L_0, the generating function (20) – by virtue of Wick theorem – will be a sum of integrals represented by Feynmann diagrams, with lines (propagators) corresponding to free Green functions (A.4), (A.5) and (23) and vertices corresponding to L_I. Suppose that $t(s)$ is varied by infinitesimal δt. Let us consider the variation of Green functions. Note that

$$\delta\langle X(t)X^*(t')\rangle_0 = (\delta t\partial_t + \delta t'\partial_{t'})\,\langle X(t)X^*(t')\rangle_0 \tag{24}$$

where $X = A, B, \psi$. The free Green function depends only on endpoint times t,t', not the shape of CTP between them.

Now, each diagram corresponds to average of the type

$$\int_C dt_1\cdots dt_N\left\langle\prod_{n\leq N}L_I(t_n)\prod_{k>N}X_k(t_k)\right\rangle_0, \tag{25}$$

where $X = \psi, A, B$ and t_k lie in the fixed (Keldysh) part of the contour for $k > N$. So we get

$$\begin{aligned}\delta\int_C dt_1\cdots dt_N\left\langle\prod_n L_I(t_n)\prod_k X_k(t_k)\right\rangle_0 &= \\ \int ds_1\cdots ds_N\sum_{m\leq N}\frac{\partial}{\partial s_m}\times \\ \left\langle\delta t_n\prod_{n\leq N}L_I(t_n)\prod_{k>N}X_k(t_k)\right\rangle_0,\end{aligned} \tag{26}$$

where we used Wick-decoupling into products of free Green functions, Leibniz rule $\delta(AB) = B\delta A + A\delta B$, $\delta dt = d\delta t = ds\partial_s\delta t$, (24), inverse Leibniz rule applied to ∂_s and back Wick-coupling. The final result of the shape variation integral is zero, which completes the proof.

IV. LORENTZ INVARIANCE

It is natural to expect that Green functions are invariant under relativistic Lorentz transformations at zero temperature if the Lagrangian is Lorentz-invariant. Such a situation should occur for vacuum state in quantum electrodynamics and generally Standard Model. However, starting from CTP makes it not obvious because the contour prefers some time direction. Intuitively, we expect that this should not bring about any problem but the warning light comes already from the non-invariance due to finite temperatures, which enters only as a jump in the contour. Below, we show that the Green function in zero-temperature vacuum are indeed invariant under continuous Lorentz transformations in Lorentz-invariant field theories but not necessarily under time reversal.

A. Relativistic notation in CTP

Let us briefly recall relativistic notation. The components of position four-vector are denoted x^μ, $\mu = 0,1,2,3$, with time $x^0 = t$ and three spatial coordinates x^k, $k = 1,2,3$. Fields, functions of position will be denoted as $f(x)$ where $x = (x^0, x^1, x^2, x^3)$. The derivatives are denoted by $\partial_\mu = \partial/\partial x^\mu$. From now on, Greek indices are reserved for four-vectors and higher Lorentz-like tensors and structures. It is extremely important for the purpose of this paper, that the time $x^0(s)$ is here a complex function of the real parameter s. In particular time derivative ∂_0 is translated into the real derivative ∂_s by

$$\partial_0 = (dx^0/ds)^{-1}\partial/\partial s \tag{27}$$

Similarly, we introduce Dirac four-delta, including complex time,

$$\delta^4(x-x') = \frac{\delta(s-s')}{dx^0/ds}\delta(x^1-x'^1)\delta(x^2-x'^2)\delta(x^3-x'^3). \tag{28}$$

and four-integral $\int_C d^4x$ with $d^4x = dx^0dx^1dx^2dx^3$ and $dx^0 = (dt/ds)ds$. For a compact notation of relative four position including the complex time component of the fourvector on CTP we will denote $|x-x'|^\mu$ with $|x-x'|^{1,2,3} = (x-x')^{1,2,3}$ but

$$|x-x'|^0 = \theta(x^0-x'^0)(x^0-x'^0) + \theta(x'^0-x^0)(x'^0-x^0), \tag{29}$$

where $\theta(x^0-x'^0) = \theta(s-s')$ applies to complex x^0 and x'^0 along CTP.

The rest of conventions, including Lorentz generators J, spinors and Lagrangian density $\mathcal{L}$ (becomes Lagrangian L when space-integrated) are consistent with textbooks [1, 38, 41], which we recall in Appendix C for completeness.

The essential part of QFT is that the Lagrangian density is a Lorentz-invariant. It means that the generating functional (20) can be written now as

$$\begin{aligned}e^{S[\chi,\xi,\eta]} \propto \int DADBD\bar{\psi}D\psi\,\exp\int_C id^4x\times \\ \left(\mathcal{L} + \sum_k(\chi_kA_k + \xi_k\cdot B_k + \bar{\eta}_k\psi_k + \bar{\psi}_k\eta_k)\right)\end{aligned} \tag{30}$$

where all fields are functions of spacetime, e.g. $\eta_k(x)$ with complex time $x^0 = t$, $\mathcal{L}$ transforms as a scalar field and normalization imposes constraint $S[0,0,0] = 0$. Here A_k denotes Lorentz scalars and B^μ_k – fourvectors with the dot $\cdot$ denoting Minkowski scalar product. In our convention, assuming that $\mathcal{L}$ is a scalar we also assumed that average



FIG. 3: Zero-temperature CTP. The boundary times t_i and t_f are sent to imaginary infinity, $i\infty$ and $-i\infty$, respectively.

charge densities are zero. The first Minkowski coordinate (time) has the jump $x^0_i - x^0_f = i\beta$.

We stress that the conservation law (B.22) extends in a straightforward way to the relativistic case, known as Ward-Takahashi identity [38]

$$\langle \partial_\mu j^\mu_N \cdots \rangle = 0,\ j^\mu_N = \sum_{n\in N} \bar{\psi}_n \gamma^\mu \psi_n. \tag{31}$$

The great advantage of Lagrangian and path integral formalism is that it seems to be perfectly Lorentz invariant at first sight. However, the possible caveats are hidden in the shape of CTP, which is bent into imaginary time but not spatial coordinates. The common sense intuition tells us that the Lorentz invariance must be broken by finite temperatures, which essentially dictate the contour jump. Moreover, even at zero temperature, the contour retains preferred time direction so Lorentz invariance is not manifest and self-evident.

B. Formal proof of invariance

We shall now present a brief formal proof of Lorentz invariance of vacuum generating function (30) at zero temperature $k_BT = 0$ or $\beta = +\infty$. Note that zero temperature makes the jump on CTP between t_i and t_f infinite. Due to the freedom of choice of the shape, we assume the contour in Fig. 2 with $t_i \to i\infty$ and $t_f \to -i\infty$, see Fig. 3. Applying a boost Λ to the fields $A \to \Lambda A$, $B \to \Lambda B$, $\psi \to \Lambda\psi$ we have the complex-time invariance of Lagrangian

$$\begin{aligned} &\int_C d^4x\mathcal{L}[A,B,\psi] \to \int d^4x\mathcal{L}(\Lambda A,\Lambda B,\Lambda\psi) = \\ &\int_C d^4x\Lambda_0\mathcal{L}[A,B,\psi] = \int_C d^4(\Lambda' x)\mathcal{L}[A,B,\psi] \\ &= \int_C d^4x\mathcal{L}[A,B,\psi] \end{aligned} \tag{32}$$

where we have used the fact that both the measure d^4x and the integration boundary is invariant under Lorentz transformations. Moreover $D(\Lambda A) = DA$, $D(\Lambda B) = DB$ and $D(\Lambda\psi) = D\psi$. All the exponent in the generating functional (30) is a Lorentz scalar so the above reasoning remains valid. We write

$$\begin{aligned} &\int DADBD\bar{\psi}D\psi\ \exp\int_C id^4x \Bigg(\mathcal{L}[A,B,\psi] + \\ &\sum_k(\Lambda\chi_k A_k + \Lambda\xi_k\cdot B_k + \Lambda\bar{\eta}_k\psi_k + \bar{\psi}_k\Lambda\eta_k)\Bigg) \\ &= \int D(\Lambda A)D(\Lambda B)D(\Lambda\bar{\psi})D(\Lambda\psi)\exp\int_C id^4x \times \\ &\Bigg(\mathcal{L}[\Lambda A,\Lambda B,\Lambda\psi] + \\ &\sum_k(\Lambda\chi_k A_k + \Lambda\xi_k\cdot B_k + \Lambda\bar{\eta}_k\Lambda\psi_k + \Lambda\bar{\psi}_k\Lambda\eta_k)\Bigg) = \\ &\int DADBD\bar{\psi}D\psi\ \exp\int_C id^4x \times \\ &\left(\mathcal{L}[A,B,\psi] + \sum_k(\chi_k A_k + \xi_k\cdot B_k + \bar{\eta}_k\psi_k + \bar{\psi}_k\eta_k)\right), \end{aligned} \tag{33}$$

which proves the invariance. However, we had to assume that Λ acts on functions of complex arguments, not just real domain, as time is complex. To be correct we should work only in the real domain (s, x^1, x^2, x^3) and define Λ by Taylor expansion of generator exponential

$$\Lambda = e^{i\omega_{\mu\nu}J^{\mu\nu}} = \sum_{n=0}^{\infty}\frac{1}{n!}(i\omega_{\mu\nu}J^{\mu\nu})^n \tag{34}$$

We noticed an analogy between the above proof and the Bohr-van Leeuwen theorem [40], that states that magnetization is absent in classical statistical physics. That theorem is proved by erasing the effect of magnetic vector potential by a shift in momentum space. Here we erase the effect of a Lorentz boost by a complex transformation of fields.

The above proof, although formally correct, is hardly acceptable because ones uses ill-defined mathematical operations. The vague issues are:

- Lorentz transformations are analytically continued to complex spacetime.
- We ignored possible problems for $x^0 \to \pm i\infty$.
- Convergence and renormalization is not discussed.

To give more insight to the proof, dispel possible objections and make it robust, we will now reformulate it in terms of perturbative diagrams showing that each diagram separately is Lorentz invariant.

C. Free theories

We shall first prove Lorentz invariance of vacuum zero-temperature Green functions in free theories, where they can be found exactly. The Lagrangian density for free scalar theory reads

$$2\mathcal{L}_0 = \partial_\mu A \partial^\mu A - m^2 A^2, \tag{35}$$

where m is the mass of the field. The Green function (propagator) satisfies Klein-Gordon equation

$$(\partial_\mu \partial^\mu + m^2)\langle A(x)A(x')\rangle_0 = -i\delta^4(x-x'), \tag{36}$$

where ∂ acts on x (we shall write ∂' if acting on x'). The solution reads

$$\begin{gathered} G_0(x,x') = \langle A(x)A(x')\rangle_0 = \\ \int \frac{d^3p}{2p^0(2\pi)^3} \sum_\pm \mp \frac{e^{\pm ip^\mu|x-x'|_\mu}}{1-e^{\pm\beta p^0}} \end{gathered} \tag{37}$$

where p^μ is the four momentum with $p^0 = \sqrt{\mathbf{p}^2+m^2}$. For the Dirac field,

$$\mathcal{L}_0 = \bar{X}(i\gamma^\mu\partial_\mu - m)X \tag{38}$$

we have

$$(-i\gamma^\mu\partial_\mu + m)_{bc}\langle X_c(x)\bar{X}_a(x')\rangle_0 = -i\delta_{ab}\delta^4(x-x'), \tag{39}$$

with the summation over c and $m_{ab} = m\delta_{ab}$. Now, we get two cases. If $X = A$ (bosonic field, unphysical, usually appearing in renormalization) then $A(t_i) = A(t_f)$ but if $X = \psi$ (fermionic field) then $\psi(t_i) = -\psi(t_f)$. The general solution reads

$$\langle X_a(x)\bar{X}_b(x')\rangle_0 = (i\gamma^\mu\partial_\mu + m)_{ab} G_0^X(x,x'), \tag{40}$$

where

$$\begin{aligned} G_0^A(x,x') &= G_0(x,x'), \\ G_0^\psi(x,x') &= \int \frac{d^3p}{2p^0(2\pi)^3} \sum_\pm \mp \frac{e^{\pm ip^\mu|x-x'|_\mu}}{1+e^{\pm\beta p^0}}. \end{aligned} \tag{41}$$

For vector fields B^μ, the internal Lorentz structure is added by $\langle B^\mu(x)B^\nu(x')\rangle_0 = ag^{\mu\nu}G_0(x,x')$. Possible problems occur for expressions of the type $\partial_\mu\partial'_\nu G_0$ because the derivative produces unphysical contact terms $\sim \delta(x^0 - x'^0)$ [41]. Fortunately, they appear always in pairs with a renormalization field, which cancels the singularity.

Nonzero temperature $\beta < \infty$ will break Lorentz invariance, which means that there is a fourvector related to temperature. There is no unique choice. One can define β^μ [29, 42] with $\beta^0 = \beta$ and $\beta^k = 0$, $k = 1,2,3$ in a particular reference frame. However, good alternatives are $k_B T^\mu = \beta^\mu/\beta\cdot\beta$ or $u^\mu = \beta^\mu/\sqrt{\beta\cdot\beta}$, see also a general discussion [43]. However, one has to remember that the zero temperature limit means *infinitely timelike* β^μ. The zero four-vector β limit, when Lorentz invariance might be obvious, corresponds unfortunately to infinite temperature so this is not what we are looking for.

In the zero-temperature limit $\beta \to +\infty$, (37) and (41) take the form

$$G_0^\psi(x,x') = G_0(x,x') = \int \frac{d^3p}{2p^0(2\pi)^3} e^{-ip^\mu|x-x'|_\mu} \tag{42}$$

Applying Lorentz generator

$$J_0 = J_{0x} + J_{0x'},\ J_{0x}^{\mu\nu} = i(x^\mu\partial^\nu - x^\nu\partial^\mu) \tag{43}$$

acting on x and x' by J_{0x} and $J_{0x'}$, respectively, one can check explicitly (a lengthy textbook exercise) that $JG = 0$ (so $\Lambda G = G$) when applied to (42), so the free theory is indeed Lorentz invariant (there are no additional problems with transformation of internal vector or spinor structure). Remember that the time derivative in J is always taken along the contour, namely with complex time derivative ∂_0 (27).

D. Interaction

The rest of the proof is similar to the shape-independence proof. The Green functions

$$\left\langle \prod_k X_k(x_k) \right\rangle \tag{44}$$

are, by virtue of Wick theorem, represented by sums of diagrams with external vertices k (not integrated), internal vertices (integrated) corresponding to the interaction part of the Lagrangian density $\mathcal{L}_I$ and connecting lines corresponding to free Green functions (37) and (41). We want to show that J vanishes when applied to the whole diagram. Suppose that the diagram has fixed number N of internal vertices $1..N$ and K external ones $N+1..N+K$. then we have to show that

$$\int d^4x_1 \cdots d^4x_N \sum_{l>N} J^l \left\langle \prod_{n\leq N} \mathcal{L}_I(x_n) \prod_{k>N} X_k(x_k) \right\rangle_0 = 0, \tag{45}$$

where X_k represent fields whose expectation value define the Green function, J^m acts on all fields meeting at the vertex m. The average is taken with respect to free Lagrangian denisty $\mathcal{L}_0$. We decouple (45) into free Green functions using Wick theorem and use Leibniz rule. Since Lorentz transformations do not affect free Green functions, we have

$$\sum_{m=1}^{N+K} J^m \left\langle \prod_{n\leq N} \mathcal{L}_I(x_n) \prod_{k>N} X_k(x_k) \right\rangle_0 = 0. \tag{46}$$

Since $\mathcal{L}_I$ is a Lorentz scalar we can change $\sum_n J^n \to \sum_n J_0^n$. Finally, we perform integration

$$\int d^4x\, J_0^{\mu\nu} \cdots \tag{47}$$

over each vertex $n \leq N$ ($x = x_n$), which yields only boundary values due to $\int dt\, x^k \partial_0 = \int ds\, x^k \partial_s$, see Eq. (C.7). However, free propagators vanish exponentially at large distances and, at zero temperature, with $x^0 \to \pm i\infty$. Hence (47) vanishes along with the left hand side of (45), which completes the proof.

This is our main result. The proof has been obtained in time-domain, without stretching the real time to infinities. Note the analogy to contour shape independence. The heart of the proof is the fact that analytic continuation of time does not hurt the generating function. It is quite intuitive but never before shown explicitly.

V. ENERGY-MOMENTUM REPRESENTATION

It is tempting to ask: Why not to prove Lorentz-invariance using simply energy-momentum representation [13, 14, 25]? This approach seems attractive and practical but it has to be supplemented by arguments following from space-time CTP representation. The main problems arise for off-shell Green functions (not describing scattering processes) and field mass/strength renormalization. They cannot be resolved without returning to space-time representation.

The energy-momentum representation can be used only on the Keldysh part of the contour stretched to the whole real axis ($t_0 \to +\infty$ and $t(s_\pm) \to -\infty$), taking separately its upper and lower branch, as Fourier transform is only defined in real axis. The Green function gets additional indices to keep track of the branch, $G_{uv}(x, x') = G(x_u, x'_v)$ where $u, v = \pm$ and $x^0_\pm = x^0 \pm i\epsilon$, $\epsilon \to 0_+$. We have

$$G(x, x') = \int \frac{d^4p}{(2\pi)^4} G(p) e^{-ip^\mu(x_\mu - x'_\mu)}, \tag{48}$$

where, in the space $(+, -)$,

$$G^\pm(p) = \begin{pmatrix} i(p^2 - m^2 + i\varepsilon)^{-1} & 2\pi\delta_-(p^2 - m^2) \\ 2\pi\delta_+(p^2 - m^2) & -i(p^2 - m^2 - i\varepsilon)^{-1} \end{pmatrix}$$
$$\pm 2\pi\delta(p^2 - m^2) n^\pm(|p^0|) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \tag{49}$$

where $\delta_\pm(p^2 - m^2) = \theta(\pm p^0)\delta(p^2 - m^2)$, the upper index $+/-$ denotes bosonic/fermionic field with $n^\pm(q) = (e^{\beta q} \mp 1)^{-1}$. We denoted $\varepsilon \to 0_+$ defined for momenta to distinguish it from ϵ defined in space-time.

In the zero-temperature limit $n \to 0$ and the free Green functions are manifestly Lorentz invariant. The problems arise for interactions. Now, the interacting Green function takes on renormalization corrections,

$$G_{0++}(p) = \frac{i}{p^2 - m^2 + i\varepsilon} \to G_{++}(p) = \frac{iZ(p)}{p^2 - M^2 + i\varepsilon}, \tag{50}$$

where $Z(p)$ is regular at $p^2 = M^2$. The renormalization is obtained by summation of Dyson series of self-energy contributions. However, such step is ill-defined for $p^2 = M^2$ because ε dominates in this regime. This can be cured by a slight tilt of the real-time contour as in Fig. 4 [38]. In fact, the new contour contains the essential feature of Fig. 3 – the endpoints in $\pm i\infty$. However, for $G_{\pm\mp}$ the trick with the contour does not help. We are forced to simply accept


$t_i \to (1 + i\varepsilon)\infty$
$s_i \to -\infty$
Im t
$\beta = +\infty$
Re t
$t_f \to (1 - i\varepsilon)\infty$
$s_f \to +\infty$

FIG. 4: The tilted Keldysh contour which circumvents the problem of pole renormalization $p^2 \sim m^2 \to M^2$

$$\delta_\pm(p^2 - m^2) \to Z\delta_\pm(p^2 - M^2) \tag{51}$$

because this part is only defined for $p^2 = M^2$. One can circumvent the problem by *ad hoc* defining extrapolation of $G_{\pm\pm}$ to $G_{\pm\mp}$ and the $p^2 = M^2$ case. Another argument is the fluctuation dissipation theorem, which imposes relations between elements of G [12], or mass derivative [33]. However, this is only the case of elementary particles like electron and photon. Composed bound states like nucleons, atoms and molecules get their renormalization corrections, too, but they cannot be simply reduced to a single dressed propagator line. In time domain, bound states do not lead to any problems because the poles/deltas in propagators appear only in the infinite time limit and we do not have to sum infinite series of diagrams.

Further problems may arise from pinch singularities [24], which may occur for products of propagators $[(p^2 - m^2 + i\varepsilon)(p^2 - m^2 - i\varepsilon)]^{-1}$. Fortunately, a careful analysis makes them always cancel [25]. Having the invariance of in-out amplitudes, it would be easy to prove Lorentz invariance if all the Green functions factorized at zero temperature [14, 29] , namely

$$\left\langle \prod X_k(p_k) \right\rangle = \left\langle \prod_{k+} X_k(p_k) \right\rangle \left\langle \prod_{k-} X_k(p_k) \right\rangle \tag{52}$$

where $k\pm$ denotes fields defined on the upper(+)/lower(−) Keldysh branch by the Fourier transform $X_\pm(p) = \int d^4x_\pm e^{ip^\mu x_{\mu\pm}}$. An attempt to prove the factorization requires generalization of CTP to the contours like that in Fig. 5 with varying σ [13, 14, 29]. If $\sigma = \beta/2$ then the factorization occurs indeed for $\beta \to \infty$. However, not every Green function is σ-independent [28, 44]. From translational invariance the nonvanishing Green functions satisfy $\sum p = 0$. The new vertical part of the contour gives additional factors $e^{\pm\sigma p^0}$ to each propagator from lower to upper Keldysh branch (+) or



FIG. 5: The separated upper and lower Keldysh contour branches $0 \leq \sigma \leq \beta$

viceversa $(-)$. These factors cancel only if

$$\sum_{k+} p_k^0 = \sum_{k-} p_k^0 = 0, \tag{53}$$

which means that the frequencies/energies of upper and lower branch *separately* add up to zero as seen in Fig. 6. This situation is common to scattering processes but not in general.

Having presented above the complications in energy-momentum space, we conclude that the proof is fully correct in space-time domain, while in the momentum/Fourier space it would require assuming at best several additional technical rules and may not be general (for very complicated diagrams). Still, most of practical calculations are easier performed in the latter case, with space-time arguments used only when the momentum rules are ambiguous.

A. Absence of time reversal symmetry

Finally, the momentum representation hides the fact that the Lorentz invariance is not necessarily valid for time reversal, which is only true for second order correlations or if the factorization (52) holds for higher order correlations. Note that the reversal $p^0 \to -p^0$ break the time symmetry because of $\theta(\pm p^0)$ factors in $G_{\pm\mp}$. Trying to reverse branches $+ \leftrightarrow -$ does not help either, because then $G_{\pm\pm}$ gets conjugated, so there is no way to restore original Green function. If we both reverse p^0 and make conjugation then we will return to the original Green function but the time will be also twice reversed which is again not what we wanted to have. The time (generally charge-parity-time) symmetry is valid only for scattering processes or spacelike separated points.

VI. CONCLUSIONS

We have shown that the closed time path formalism is consistent with Lorentz invariance at zero temperature. The presented picture of dual operator-Hamiltonian and path-integral-Lagrangian fits well into covariant relativistic framework with the analytic continuation of time as the only nontrivial extra element. It is remarkable that complex time glues statistical mechanics, time evolution and Lorentz symmetry without any problems. The fact that CTP is invariant only under continuous transformations, not time reversal, shows that there the properties of CTP still differ from scattering amplitudes. An open issue is whether the Lorentz symmetry can be shown non-perturbatively (what about spontaneous breaking?) and for quantum gravity.



FIG. 6: The particle view of factorization mechanism. The Green function is σ-independent only if $p_{1+}^0 + p_{2+}^0 + p_{3+}^0 + p_{4+}^0 + p_{5+}^0 = 0$ and $p_{6-}^0 + p_{7-}^0 + p_{8-}^0 = 0$

Acknowledgements

I thank W. Belzig, A. Neumaier, S. Mrówczyński and P. Chankowski for fruitful discussions and motivation and for hospitality of University of Konstanz, where part of this work has been completed.

Appendix A

(a). Free Green functions

The most convenient free Hamiltonian is the quadratic form of bosonic and fermionic operators

$$\hat{H}_0 = \sum_{kl} (b_{kl}\hat{x}_k\hat{x}_l + f_{kl}\hat{\phi}_k\hat{\phi}_l) \tag{A.1}$$

where $\hat{x}$ and $\hat{\phi}$ are Hermitian operators with bosonic and fermionic commutation relations, respectively, $[\hat{x}_k, \hat{x}_l] = ig_{kl}\hat{1}$, $\{\phi_k, \phi_l\} = h_{kl}\hat{1}$, with real g and h.

One can always diagonalize $\hat{H}_0$ so that

$$\hat{H}_0 = H_{vac} + \sum_k E_k(\hat{A}_k^\dagger \hat{A}_k + \hat{\psi}_k^\dagger \hat{\psi}_k) \tag{A.2}$$

where H_{vac} is the vacuum energy (can be ignored), $\hat{A}$ and $\hat{\psi}$ are linear combinations of $\hat{x}$ and $\hat{\phi}$, respectively, with the property

$$\begin{aligned} &[\hat{A}_k, \hat{A}_l] = 0,\ [\hat{A}_k, \hat{A}_l^\dagger] = \delta_{kl}, \\ &\{\hat{\psi}_k, \hat{\psi}_l\} = 0,\ \{\hat{\psi}_k, \hat{\psi}_l^\dagger\} = \delta_{kl}. \end{aligned} \tag{A.3}$$

It is especially simple and instructive to find Green functions for the above Hamiltonian, according to (12) and (16), extended to the whole CTP. In the case of bosonic operators, $\langle A(t)\rangle_0 = \langle A^*(t)\rangle_0 = 0$ and

$$\begin{aligned}
&\langle A_k(t)A_l(t')\rangle_0 = \langle A^*_k(t)A^*_l(t')\rangle_0 = 0,\\
&\langle A_k(t)A^*_l(t')\rangle_0 = \delta_{kl}e^{i(t'-t)E_k}\times\\
&\left[\frac{\theta(t-t')}{1-e^{-\beta E_k}}+\frac{\theta(t'-t)}{e^{\beta E_k}-1}\right]. \qquad (A.4)
\end{aligned}$$

For fermionic operators, it is important that ϕ (ψ, ψ^*) is never a c-number but a Grassmann number with the property $\phi_a\phi_b = -\phi_b\phi_a$, which follows from anticommutation relations. The anticommutation makes it necessary to use different boundary conditions, due to Kubo, Martin and Schwinger [35], imposing the sign reversal on the jump $t_f \to t_i$, so the Green functions read $\langle\psi(t)\rangle_0 = \langle\psi^*(t)\rangle_0 = 0$ and

$$\begin{aligned}
&\langle \psi_k(t)\psi_l(t')\rangle_0 = \langle \psi^*(t)_k\psi^*_l(t')\rangle_0 = 0,\\
&\langle \psi_k(t)\psi^*_l(t')\rangle_0 = \delta_{kl}e^{i(t'-t)E_k}\times\\
&\left[\frac{\theta(t-t')}{1+e^{-\beta E_k}}-\frac{\theta(t'-t)}{e^{\beta E_k}+1}\right]. \qquad (A.5)
\end{aligned}$$

In both cases, the Green functions satisfy the equation

$$\begin{aligned}
&\delta(t-t')\delta_{kl} = [\partial_t + iE_k]\langle A_k(t)A^*_l(t')\rangle_0\\
&= [\partial_t + iE_k]\langle \psi_k(t)\psi^*_l(t')\rangle_0 \qquad (A.6)
\end{aligned}$$

with different boundary conditions, $A(t_i) = A(t_f)$ but $\psi(t_i) = -\psi(t_f)$.

It is often convenient to introduce the Keldysh space for Re $t > 0$ in analogy to superoperators (10). In particular $x^\pm(t) = x(t_\pm)$, $\phi^\pm(t) = \phi(t_\pm)$. and similarly $2x^c(t) = x^+(t) + x^-$ ($2\phi^c(t) = \phi^+(t) + \phi^-(t)$) and $x^q(t) = x^+(t) - x^-(t)$ ($\phi^q(t) = \phi^+(t) - \phi^-(t)$). From (13) it follows that

$$\langle x^q_k(t)x_l(t')\rangle_0 = \langle \phi^q_k(t)\phi_l(t')\rangle_0 = 0 \qquad (A.7)$$

for $\mathrm{Re}\,t' < t$. The general structure of Keldysh Green functions reads

$$\begin{aligned}
&\langle x^q_k(t)x^q_l(t')\rangle_0 = \langle \phi^q_k(t)\phi^q_l(t')\rangle_0 = 0,\\
&\langle A^c_k(t)A^{c*}_k(t')\rangle_0 = e^{i(t'-t)E_k}/2\tanh(\beta E_k/2),\\
&\langle \psi^c_k(t)\psi^{c*}_k(t')\rangle_0 = e^{i(t'-t)E_k}/2\coth(\beta E_k/2),\\
&\langle A^c_k(t)A^{q*}_k(t')\rangle_0 = \langle \psi^c_k(t)\psi^{q*}_k(t')\rangle_0 \qquad (A.8)\\
&= e^{i(t'-t)E_k}\theta(t-t'),\\
&\langle A^q_k(t)A^{c*}_k(t')\rangle_0 = \langle \psi^q_k(t)\psi^{c*}_k(t')\rangle_0\\
&= -e^{i(t'-t)E_k}\theta(t'-t).
\end{aligned}$$

In the above equations t is the ordinary variable (we forget temporarily about s) and θ is the usual Heaviside function. Note that the last two equations indicate causality – q-variables influence the future but not the past.

(b). Wick theorem

The many-point Green functions are obtained from Wick theorem [36, 37]. For products of odd number of operators, Green function vanishes while for the even number $2n$,

$$\begin{aligned}
&\left\langle \prod_{k=1}^{2n} x_k(t_k)\right\rangle_0 =\\
&\frac{1}{n!2^n}\sum_\sigma\prod_{k=1}^{n}\langle x_{\sigma(2k-1)}(t_{\sigma(2k-1)})x_{\sigma(2k)}(t_{\sigma(2k)})\rangle_0,\\
&\left\langle \prod_{k=1}^{2n} \phi_k(t_k)\right\rangle_0 = \qquad (A.9)\\
&\frac{\mathrm{sgn}\,\sigma}{n!2^n}\sum_\sigma\prod_{k=1}^{n}\langle \phi_{\sigma(2k-1)}(t_{\sigma(2k-1)})\phi_{\sigma(2k)}(t_{\sigma(2k)})\rangle_0
\end{aligned}$$

The Wick theorem states essentially that every many-point Green function for a quadratic Hamiltonian splits into products of two-point functions. This fact is analogous to the property of Fourier transforms of Gaussian functions, which are again Gaussian. We will return to this interpretation in Appendix B(c). Wick theorem is useful for interactions – one proceeds perturbatively, expanding (18) in interaction strength.

Appendix B

(a). Bosonic fields

Let us begin with the Lagrangian

$$L(q,\dot q,t) = \sum_k (m_k\dot q_k^2/2 + f_k(q)\dot q_k) - g(q) \qquad (B.1)$$

for generalized coordinates q and $\dot q = dq/dt$. The classical Hamiltonian is obtained by Legendre transformation, defining momenta,

$$p_k = \partial L/\partial \dot q_k = m_k\dot q_k + f_k(q) \qquad (B.2)$$

so that the Hamiltonian reads

$$H(q,p) = \sum_k p_k\dot q_k - L = \sum_k (p_k - f_k(q))^2/2m_k + g(q). \qquad (B.3)$$

We assume quantum evolution of the wavefunction $\Psi(q)$ given by the path integral

$$\Psi(\tilde q,t) = \int_{q(0)=q'}^{q(t)=\tilde q} Dq\, e^{\int iL(q,\dot q)dt}\Psi(q',0), \qquad (B.4)$$

where Dq includes all necessary normalization factors. To find the operator form of the Hamiltonian, we expand the above equation for small times,

$$\Psi(\tilde{q},t)=\int^{q(t)=\tilde{q}} Dq\, e^{\int_0^t idt'\sum_k m_k\dot{q}_k^2(t')/2}\times$$
$$\Bigg(1-g(\tilde{q})t+\sum_k\int_0^t idt'\dot{q}_k(t')f_k(\tilde{q})$$
$$-\int_0^t dt'\dot{q}_k(t')\frac{\partial}{\partial\tilde{q}_k}+\sum_{kl}\int_0^t dt'dt''\dot{q}_k(t')\dot{q}_l(t'') \tag{B.5}$$
$$\left(\frac{\partial^2}{2\partial\tilde{q}_k\partial\tilde{q}_l}-i\partial f_k(\tilde{q})\frac{\partial}{\partial\tilde{q}_l}-f_k(\tilde{q})f_l(\tilde{q})/2\right)$$
$$-\sum_{kl}\int_0^t idt'\dot{q}_k(t')\int_0^{t'}dt''\dot{q}_l(t'')\frac{\partial f_k(\tilde{q})}{\partial\tilde{q}_l}\Bigg)\Psi(\tilde{q},0).$$

The ignored terms are of the order $\sim t^2$. We now use the fact that

$$\int Dq\, e^{\int idt'\sum_k m_k\dot{q}_k^2(t')/2}\dot{q}_k(t')=0,$$
$$\int Dq\, e^{\int idt'\sum_k m_k\dot{q}_k^2(t')/2}\dot{q}_k(t')\dot{q}_l(t'') \tag{B.6}$$
$$=\frac{i}{m_k}\delta_{kl}\delta(t'-t''),$$

which gives

$$(\Psi(\tilde{q},t)-\Psi(\tilde{q},0))/it=\Bigg(-g(\tilde{q})+\sum_k\frac{1}{m_k}\times \tag{B.7}$$
$$\left(\frac{\partial^2}{2\partial\tilde{q}_k^2}-f_k(\tilde{q})\frac{i\partial}{\partial\tilde{q}_k}-f_k^2(\tilde{q})/2-\frac{i\partial f_k(\tilde{q})}{2\partial\tilde{q}_k}\right)\Bigg)\Psi(\tilde{q},0).$$

Let us shortly comment the last term in bracket. It has been obtained by assuming $\theta(t)\delta(t)\to\delta(t)/2$ or symmetrizing the last integral in (B.5) [39]. However, in the fundamental theories, like quantum electrodynamics, weak, strong interactions and generally Standard Model this term is absent. It follows from the fact that the even the most dangerous terms, due to non-Abelian gauge fields, have $\partial f_k/\partial q_k=0$. Nevertheless, this indicates potential problems for completely general Lagrangians, especially for gravity.

Finally, we get

$$i\partial_t\Psi=\hat{H}\Psi \tag{B.8}$$

where

$$\hat{H}=\sum_k(\hat{p}_k-f_k(q))^2/2m_k+g(q),\ \hat{p}_k=\frac{\partial}{i\partial q_k}, \tag{B.9}$$

which proves the equivalence between bosonic path integrals and Hamiltonian picture, with the commutation rule $[\hat{q}_k,\hat{p}_l]=i\delta_{kl}$.

(b). Fermionic fields

Fermionic path integrals are more complicated due to anticommutation rules. We have to define abstract anticommuting Grassmann numbers, ϕ_k, with the property $\phi_k\phi_l=-\phi_l\phi_k$ and the integrals,

$$\int d\phi_k=0,\ \int d\phi_k\,\phi_k=1, \tag{B.10}$$
$$\int d\phi_1\cdots d\phi_n\,\phi_n\cdots\phi_1=1. \tag{B.11}$$

One can check that

$$\int D\phi^*D\phi e^{-\sum_{kl}\phi_k^*F_{kl}\phi_l}=\det F, \tag{B.12}$$
$$\int D\phi^*D\phi e^{-\sum_{kl}\phi_k^*F_{kl}\phi_l}\phi_j\phi_m^*=(F^{-1})_{jm}\det F.$$

Here $\phi=\phi_r+i\phi_i$ and $\phi^*=\phi_r-i\phi_i$ are independent fields and $d\phi^*d\phi=id\phi_id\phi_r/2$.

Let us now take an abstract Lagrangian

$$L_0(\psi,\dot{\psi},t)=\sum_k\psi_k^*(i\dot{\psi}_k-E_k\psi_k). \tag{B.13}$$

According to the classical rule we can construct an abstract Hamiltonian

$$H_0=\sum_k p_k\dot{\psi}_k-L_0=\sum_k E_k\pi_k\psi_k \tag{B.14}$$

for

$$\pi_k=\frac{\partial L_0}{\partial\psi_k}=i\psi_k^*. \tag{B.15}$$

In the continuous case for CTP,

$$\frac{\int D\psi^*D\psi e^{\sum_n\int_C idt\psi_n^*(i\partial_t-E_n)\psi_n}\psi_k(\tilde{t})\psi_l^*(t')}{\int D\psi^*D\psi e^{\sum_n\int_C idt\psi_n^*(i\partial_t-E_n)\psi_n}} \tag{B.16}$$
$$=\delta_{kl}e^{i(t'-\tilde{t})E_k}\left[\frac{\theta(\tilde{t}-t')}{1+e^{-E_k/k_BT}}-\frac{\theta(t'-\tilde{t})}{e^{E_k/k_BT}+1}\right]$$

which is the equivalent to (A.5). The equivalence between bosonic-fermionic path integrals and Hamiltonian operator can be now easily extended to general family of Lagrangians of the type

$$L=\sum_k(m_k\dot{q}_k^2/2+\dot{q}_kf_k(q,\psi)+i\psi_k^*\dot{\psi}_k)-g(q,\psi), \tag{B.17}$$

where $f_k(q,\psi)$ and $g(q,\psi)$ are real-valued and even in ψ, and Hamiltonian

$$\hat{H}=\sum_k(\hat{p}_k-f_k(\hat{q},\hat{\psi}))^2/2m_k+g(\hat{q},\hat{\psi}) \tag{B.18}$$

with commutation relations $[\hat{p}_k,\hat{q}_l]=i$ and $\{\hat{\psi}_k,\hat{\psi}_l^*\}=\delta_{kl}$. We emphasize however that the equivalence does not generalize to completely arbitrary Lagrangians, containing higher powers of time derivatives, e.g. $\dot{q}^4$ or $\dot{\psi}^2$.

(c). Wick theorem

The Wick theorem (Appendix A(b)) is now a straightforward consequence of Gaussian integrals

$$\int DqD\phi e^{-\sum_{kl}(q_k g_{kl} q_l-\phi_k b_{kl}\phi_l)/2+\sum_k(\chi_k q_k+\eta_k\phi_k)}$$
$$\propto e^{\sum_{kl}(\chi_k g^{-1}_{kl}\chi_l-\eta_k b^{-1}_{kl}\eta_l)/2}. \tag{B.19}$$

where g and b are real symmetric and antisymmetric matrices, respectively, while g^{-1} and b^{-1} are their inverses.

(d). Conservation laws

As noticed already by Kadanoff and Baym, the CTP technique maintains classical conservation laws. Suppose that the fermionic terms in Lagrangian (B.17) appear in separate sets,

$$L=\sum_{n\in N}\psi^*_n i\partial_t\psi_n+L_I, \tag{B.20}$$

where L_I is built from bosonic field and fermionic quadratic forms

$$\sum_N\sum_{n,n'\in N}\psi^*_n\psi_{n'}. \tag{B.21}$$

Sets N form a family of *disjoint* sets and L_I may contains arbitrary products and sums of (B.21) for different Ns together with bosonic fields. It is however forbidden to include terms like $\psi^*_k\psi_{k'}$ where k and k' belong to different sets N and N', respectively. Each set N defines one conserved quantity, $Q_N(t)=\sum_{n\in N}\psi^*_n(t)\psi_n(t)$. The conservation law states that

$$\langle\partial_t Q_N(t)\cdots\rangle=0, \tag{B.22}$$

when averaging with the path integral (20). To prove (B.22) we make the transformation $\psi_n(t)=e^{i\alpha(t)}\psi'_n(t)$, for real α with boundary condition $\alpha(t_i)=\alpha(t_f)$. As the transformation is linear, $D\psi^*D\psi=D\psi'^*D\psi'$ and

$$DAD\psi^*D\psi\, e^{\int_C idt\, L}=DAD\psi^*D\psi\, e^{\int_C idt\,(L(t)-Q_N\partial_t\alpha)}$$
$$=DAD\psi^*D\psi\, e^{\int_C idt\,(L(t)+\alpha\partial_t Q_N)} \tag{B.23}$$

where we performed integration by parts with respect to time in the last line. Taking the above equation integrated with arbitrary Q_N-conserving field function and preforming the functional derivative

$$\delta/\delta\alpha(t)|_{\alpha=0} \tag{B.24}$$

we obtain (B.22). The Q_N-conserving functions must be built of Hermitian forms of type (B.21) so that they are invariant under $\psi\to\psi'$. We could certainly repeat the proof perturbatively, as we did to prove shape independence (some issues are resolved by renormalization).

(e). Hubbard-Stratonovich transformation and renormalization

Here we discuss the two techinques often used in context of path integrals. The Hubbard-Stratonovich transformation [45] allows to reduce order of terms in path integral by an auxiliary bosonic field $X(t)$,

$$e^{\int idt Y^2(t)/4a}\propto\int DXe^{\int idt(\sqrt{a}X(t)Y(t)-X^2(t))}. \tag{B.25}$$

Note that $\sqrt{a}$ is *imaginary* if $a<0$ and in this case X does not correspond to any physical field although path integrals can be still performed. The renormalization is used to kill divergences appearing in quantum field theory [1]. It requires introducing unphysical fields, too. The most common form of renormalizing terms in Largangian is

$$L_R=f(X_k,i\tilde{X}_m) \tag{B.26}$$

where $\tilde{X}_m$ is the unphysical field as it appears with i. Both techniques are fully consistent with CTP formalism. For details of these techniques we refer readers to textbooks [1, 38, 41].

Appendix C

For general multi-indexed relativistic structures we adopt Einstein convention $A\cdot B=A^\mu B_\mu=\sum_{\mu=0,1,2,3}A^\mu B_\mu$ and $A^2=A\cdot A$. We denote Minkowski metric tensor $g^{\mu\nu}=g_{\mu\nu}$ equal $+1$ for $\mu=\nu=0$, -1 for $\mu=\nu=1,2,3$ and 0 otherwise. Next, g is used to lower/raise index, $A_\mu=g_{\mu\nu}A^\nu$, $A^\mu=g^{\mu\nu}A_\nu$. In this notation $g^\mu_\nu=\delta^\mu_\nu$ is equal $+1$ if $\mu=\nu$ and 0 otherwise and $A^0=A_0$ while $A^{1,2,3}=-A_{1,2,3}$. The four-vector measure is denoted $d^4x=dx^0dx^1dx^2dx^3$. We extract the usual three-vectors $\mathbf{x}=(x^1,x^2,x^3)$, its square $\mathbf{x}^2=(x^1)^2+(x^2)^2+(x^3)^3$ and measure $d^3x=dx^1dx^2dx^3$. The natural Lorentz transformation is defined on four-vectors,

$$A'^\mu=\Lambda'^\mu{}_\nu A^\nu. \tag{C.1}$$

The transformation must satisfy conservation of metric tensor

$$g^{\mu\nu}=\Lambda'^\mu{}_\alpha\Lambda'^\nu{}_\beta g^{\alpha\beta}. \tag{C.2}$$

In general, the continuous Lorentz transformation can be written in the form

$$\Lambda=\exp(-i\omega_{\mu\nu}J^{\mu\nu}/2) \tag{C.3}$$

where J is the generator of the Lorentz group, satisfying

$$[J^{\mu\nu},J^{\alpha\beta}]=i(g^{\nu\alpha}J^{\mu\beta}-g^{\mu\alpha}J^{\nu\beta}-g^{\nu\beta}J^{\mu\alpha}+g^{\mu\beta}J^{\nu\alpha}) \tag{C.4}$$

which yields $J^{\mu\nu} = -J^{\nu\mu}$. The generator for (C.1) reads

$$(J'^{\mu\nu})^{\alpha}{}_{\beta} = i(g^{\mu\alpha}g^{\nu}_{\beta} - g^{\nu\alpha}g^{\mu}_{\alpha}) \tag{C.5}$$

For the scalar field

$$\Lambda_0 A(x) = A(\Lambda'^{-1}x) \tag{C.6}$$

with the corresponding generator

$$J^{\mu\nu}_0 = i(x^\mu\partial^\nu - x^\nu\partial^\mu) \tag{C.7}$$

If the field has vector structure, e.g. $A_\mu(x)$ then the generator reads $J' + J_0$, acting both on argument x and four-vector (indices μ). The scalar fields are examples of the general family of Lorentz scalars, combinations of fields that transform according to Λ_0. Other examples are $A^\mu(x)B_\mu(x)$, $A(x)\partial_\mu B^\mu(x)$ and $\partial^\mu A\partial_\mu B$ and their combinations (products, sums, multi-indexed forms). In general, if all spacetime indices (μ) appear in pairs, then we have a Lorentz scalar.

Apart from continuous transformations there exist two special discrete transformations, time reversal $x^0 \to -x^0$ and parity (mirror) inversion $x^k \to -x^k$, $k = 1, 2, 3$. One can also make charge conjugation $e \to -e$ (the interaction strength). The proof of Lorentz invariance of zero temperature vacuum will be valid only for continuous transformations. If Lagrangian is invariant with respect to parity inversion or charge conjugation then we can include it, too (which is e.g. not the case for weak interactions).

Fermionic fields are represented by four-component spinors ψ_a, $a = 1, 2, 3, 4$, which have nothing to do with four-vectors. To define corresponding Lorentz transformations, we first denote Dirac 4×4 matrices, γ^μ, with the property

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} \tag{C.8}$$

and $\gamma^0 = \gamma^{0\dagger}$, $\gamma^k = -\gamma^{k\dagger}$, $k = 1, 2, 3$. The Lorentz generator reads,

$$J^{\mu\nu}_s = i[\gamma^\mu, \gamma^\nu]/4. \tag{C.9}$$

The Dirac matrices transform as four-vectors,

$$\Lambda^{-1}_s \gamma^\mu \Lambda_s = \Lambda'^{\mu}_{\nu}\gamma^\nu. \tag{C.10}$$

The conjugate spinor, $\psi^\dagger$ transforms as $\psi^\dagger\Lambda^\dagger_s$. Unfortunately, J_s is not Hermitian and $\Lambda^\dagger_s\Lambda_s \neq 1$ but $\Lambda^\dagger_s\gamma^0\Lambda_s = \gamma_0$ or $J^{\mu\nu\dagger}_s\gamma^0 = \gamma^0 J^{\mu\nu}_s$. Therefore we denote $\bar\psi = \psi^\dagger\gamma^0$ and $\bar\psi\phi$ is again a Lorentz scalar as well as $A_\mu\bar\psi\gamma^\mu\psi$, $\bar\psi\gamma^\mu\partial_\mu\phi$ and so on. General rules to construct a Lorentz scalar are:

- a product/sum of Lorentz scalars is a Lorentz scalar,
- $\bar\psi\phi$ and A are Lorentz scalars,
- $\bar\psi\gamma^\nu\phi$ and A^ν and $\partial^\nu A$ are Lorentz vectors,
- $\bar\psi\gamma^\nu\gamma^\mu\phi$ and $\partial^\mu A^\nu$ are Lorentz tensors, etc.,
- A product of Lorentz vectors (tensors) is a Lorentz scalar if the Greek indices always come in raised/lowered pairs ${}^{\mu}_{\mu}$ and Einstein summation is performed.

Instead of Lagrangian, in relativistic spacetime we will rather use its density $\mathcal{L}$ with the property

$$L = \int d^3x\, \mathcal{L}, \tag{C.11}$$

where $d^3x = dx^1dx^2dx^3$ is the three dimensional volume measure. It allows to write action in the spacetime

$$\int dt\, L = \int d^4x\, \mathcal{L}. \tag{C.12}$$

Every Lorentz transformation Λ conserves measure $\Lambda(d^4x) = d^4x$.

In the Lagrangian density (and other functionals) one can find every kind of combination of fields, $A(x)$, $B^\mu(x)$, ψ, etc. The Lagrangian will be Lorentz-invariant if its density transforms as Lorentz scalar

$$\mathcal{L}(\Lambda A, \Lambda B, \Lambda\psi) = \Lambda_0\mathcal{L}(A, B, \psi) \tag{C.13}$$

the generic form of such Lagrangian reads

$$\begin{aligned}
\mathcal{L} = {} & \sum_{kl}(a^0_{kl}\bar\psi_k i\gamma^\mu\partial_\mu\psi_l + a^1_{kl}(\partial_\mu A_k)(\partial^\mu A_l) \\
& + a^2_{kl}(\partial_\mu B^\mu_k)(\partial_\nu B^\nu_l) + a^3_{kl}(\partial_\mu B^\nu_k)(\partial^\mu B_{\nu l}) + \\
& + a^4_{kl}A_kA_l + a^5_{kl}B^\mu_k B_{\mu l} + a^6_{kl}\bar\psi_k\psi_l + a^7_{kl}B^\mu_k\partial_\mu A_l) \\
& + \sum_{klm}(a^8_{klm}A_m\bar\psi_l\psi_m + a^9_{klm}B_{\mu k}\bar\psi_l\gamma^\mu\psi_m + \\
& a^{10}_{klm}B^\mu_k B^\nu_l\partial_\mu B_{\nu m} + a^{11}_{klm}\partial_\mu A_k\bar\psi_l\gamma^\mu\psi_m) \\
& + \sum_{klmn}(a^{12}_{klmn}\bar\psi_k\psi_l\bar\psi_m\psi_n + a^{13}_{klmn}B^\mu_k B_{\mu l}B^\nu_k B_{l\nu} + \ldots.
\end{aligned} \tag{C.14}$$

When considering weak interactions, one should include also pseudoscalars $\bar\psi_k\gamma^5\psi_l$, $A_\mu\bar\psi_k\gamma^5\gamma^\mu\psi_l$ where $\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3$, which transforms as scalar under continuous transformations but changes sign under parity inversion.

---

[1] L.H. Ryder, *Quantum Field Theory* (Cambridge University Press, 1985); N. Bogoliubov, A.A. Logunov, A.I. Oksak, I.T. Todorov, *General Principles of Quantum Field Theory*, (Kluwer Academic Publishers, Dordrecht, 1990).

[2] S. Weinberg, *The Quantum Theory of Fields*, (Cambridge University Press, Cambridge, 1995).

[3] R.P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, New York,1965)

[4] T. Matsubara, Prog. Theor. Phys., **14**, 351 (1955).

[5] J. I. Kapusta and C. Gale, *Finite-Temperature Field Theory*, (Cambridge University Press, Cambridge, 2005)

[6] J. Schwinger, J. Math. Phys. **2**, 407 (1961).

[7] L.P. Kadanoff and G. Baym, *Quantum Statistical Mechanics*, (W.A. Benjamin, New York, 1962).

[8] L. Keldysh, Sov. Phys. JETP **20**, 1018 (1965).

[9] K. T. Mahanthappa, Phys. Rev. **126**, 329 (1962); P. M. Bakshi and K. T. Mahanthappa, J. Math. Phys. **4**, 1, 12 (1963); R. A. Craig, J. Math. Phys. **9**, 605 (1968).

[10] R. Mills, *Propagators for many-particle systems*, (Gordon and Breach, New York, 1969).

[11] R.E. Cutkosky, J. Math. Phys. **1**, 429 (1960).

[12] K. Chou, Z. Su, B. Hao and L. Yu, Phys. Rep. **118**, 1 (1985).

[13] A.J. Niemi, G.W. Semenoff, Nucl. Phys. B **220**, 181 (1984); Ann. Phys. **152**, 105 (1984).

[14] N. P. Landsman and Ch. G. van Weert, Phys. Rep. **145**, 141 (1987).

[15] R. L. Kobes, G. W. Semenoff and N. Weiss, Z. f. Physik C, **29**, 371, (1985).

[16] H. B. Callen and T. A. Welton, Phys. Rev. **83**, 34 (1951).

[17] A. Kamenev, A. Levchenko, Adv. in Phys. **58**, 197 (2009).

[18] A. M. Rey, B. L. Hu, E. Calzetta, A. Roura, and C. W. Clark, Phys. Rev. A **69**, 033610 (2004).

[19] E. Calzetta and B. L. Hu, Phys. Rev. D **35**, 495 (1987); **37**, 2878 (1988).

[20] S. Weinberg, Phys. Rev. D **72**, 043514 (2005).

[21] R. L. Kobes, K. L. Kowalski, Phys. Rev. D **34**, 513 (1986).

[22] U. Kraemmer and A. Rebhan, Rep. Prog. Phys. **67**, 351 (2004).

[23] J. Schwinger, Phys. Rev. **82**, 914 (1951).

[24] L. Dolan, R. Jackiw, Phys. Rev. D **9**, 3320 (1974).

[25] R. L. Kobes and G. W. Semenoff, Nucl. Phys. **B272**, 329 (1986); **B260**, 714 (1985).

[26] F. Gelis, Phys. Lett. B **455**, 205 (1999); Z. Phys. C, **70**, 321 (1996).

[27] M. Le Bellac and H. Mabilat, Phys. Lett. B, **381**, 262 (1996); H. Mabilat, Z. Phys. C, **75**, 155 (1997).

[28] T. S. Evans and A. C. Pearson, Phys. Rev. D **52**, 4652 (1995).

[29] R. J. Furnstahl and B. D. Serot, Phys. Rev. C **44**, 2141 (1991).

[30] V. A. Kostelecky, Phys. Rev. D **69**, 105009 (2004).

[31] R. F. Streater and A. S. Wightman *PCT, spin and statistics, and all that* (Benjamin, New York, 1964).

[32] J. von Neumann, *Mathematical Foundations of Quantum Mechanics* (Princeton U.P., Princeton, 1932).

[33] A. Niegawa, Phys. Rev. D **40**, 1199 (1989) .

[34] T. S. Evans, Phys. Rev. D **47**, R4196 (1993).

[35] R. Kubo, J. Phys. Soc. Jpn. **12**, 570 (1957); Rep. Prog. Phys. **29**, 255 (1966); P. C. Martin and J. Schwinger, Phys. Rev. **115**, 1432 (1959).

[36] G.C. Wick, Phys. Rev. **80**, 268 (1950).

[37] T. S. Evans, D. A. Steer, Nucl. Phys. **B474**, 481 (1996).

[38] M. Peskin, D. Schroeder, *An Introduction to Quantum Field Theory*, (Perseus Books, Reading, 1995).

[39] J. Zinn-Justin, *Path Integrals in Quantum Mechanics* (Oxford University Press, Oxford, 2005).

[40] II. The Doctor's Dissertation (Text and Translation), in: L. Rosenfeld and J. R. Nielsen, Editor(s), Niels Bohr Collected Works, Elsevier, 1972, vol. 1 pp. 163, 1972, 165393. http://dx.doi.org/10.1016/S1876-0503(08)70015-X; J. H. Van Leeuwen, Journal de Physique et le Radium 2 (12) 361377 (1921), http://dx.doi.org/10.1051/jphysrad:01921002012036100.

[41] C. Itzykson, J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).

[42] H.A. Weldon, Phys. Rev. D **26**, 1394 (1982).

[43] J. Dunkel, P. Hänggi, and S. Hilbert, Nature Phys. **5**, 741 (2009).

[44] M. Inui, H. Kohyama, and A. Niegawa, Phys. Rev. D **73**, 047702 (2006).

[45] R. L. Stratonovich, Sov. Phys. Dokl. **2**, 416 (1958). J. Hubbard, Phys. Rev. Lett. **3**, 77 (1959).